

COLUMBIA-SUICIDE SEVERITY RATING SCALE (C-SSRS)

Very Young Child/Cognitively Impaired – Lifetime Recent

Version 6/23/10 m4/3/18

*Posner, K.; Brent, D.; Lucas, C.; Gould, M.; Stanley, B.; Brown, G.; Fisher, P.; Zelazny, J.;
Burke, A.; Oquendo, M.; Mann, J.*

Disclaimer:

This scale is intended to be used by individuals who have received training in its administration. The questions contained in the Columbia-Suicide Severity Rating Scale are suggested probes. Ultimately, the determination of the presence of suicidal ideation or behavior depends on the judgment of the individual administering the scale.

*Definitions of behavioral suicidal events in this scale are based on those used in **The Columbia Suicide History Form**, developed by John Mann, MD and Maria Oquendo, MD, Conte Center for the Neuroscience of Mental Disorders (CCNMD), New York State Psychiatric Institute, 1051 Riverside Drive, New York, NY, 10032. (Oquendo M. A., Halberstam B. & Mann J. J., Risk factors for suicidal behavior: utility and limitations of research instruments. In M.B. First [Ed.] Standardized Evaluation in Clinical Practice, pp. 103 -130, 2003.)*

For reprints of the C-SSRS contact Kelly Posner, Ph.D., New York State Psychiatric Institute, 1051 Riverside Drive, New York, New York, 10032; inquiries and training requirements contact posnerk@nyspi.columbia.edu

SUICIDAL BEHAVIOR <i>(Check all that apply, so long as these are separate events; must ask about all types)</i>		Lifetime	Past 3 Months
Actual Attempt: A potentially self-injurious act committed with at least some wish to die, <i>as a result of act</i> . Behavior was in part thought of as method to kill oneself. Intent does not have to be 100%. If there is any intent/desire to die associated with the act, then it can be considered an actual suicide attempt. There does not have to be any injury or harm , just the potential for injury or harm. If person pulls trigger while gun is in mouth but gun is broken so no injury results, this is considered an attempt. Inferring Intent: Even if an individual denies intent/wish to die, it may be inferred clinically from the behavior or circumstances. For example, a highly lethal act that is clearly not an accident so no other intent but suicide can be inferred (e.g., gunshot to head, jumping from window of a high floor/story). Also, if someone denies intent to die, but they thought that what they did could be lethal, intent may be inferred. Did you ever <u>do anything</u> to try to kill yourself or make yourself not alive anymore? What did you do? Did you ever hurt yourself on purpose? Why did you do that? Did you _____ as a way to end your life? Did you want to die (even a little) when you _____? Were you trying to make yourself not alive anymore when you _____? Or did you think it was possible you could have died from _____? Or did you do it purely for other reasons, <u>not at all</u> to end your life or kill yourself (like to make yourself feel better, or get something else to happen)? (Self-Injurious Behavior without suicidal intent) If yes, describe:		Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of Attempts _____	Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of Attempts _____
Has subject engaged in Non-Suicidal Self-Injurious Behavior?		Yes No <input type="checkbox"/> <input type="checkbox"/> Yes No	Yes No <input type="checkbox"/> <input type="checkbox"/> Yes No
Has subject engaged in Self-Injurious Behavior, intent unknown?		<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>
Interrupted Attempt: When the person is interrupted (by an outside circumstance) from starting the potentially self-injurious act (<i>if not for that, actual attempt would have occurred</i>). Overdose: Person has pills in hand but is stopped from ingesting. Once they ingest any pills, this becomes an attempt rather than an interrupted attempt. Shooting: Person has gun pointed toward self, gun is taken away by someone else, or is somehow prevented from pulling trigger. Once they pull the trigger, even if the gun fails to fire, it is an attempt. Jumping: Person is poised to jump, is grabbed and taken down from ledge. Hanging: Person has noose around neck but has not yet started to hang - is stopped from doing so. Has there been a time when you started to do something to make yourself not alive anymore (end your life or kill yourself) but someone or something stopped you before you actually did anything? What did you do? If yes, describe:		Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of interrupted _____	Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of interrupted _____
Aborted or Self-Interrupted Attempt: When person begins to take steps toward making a suicide attempt, but stops themselves before they actually have engaged in any self-destructive behavior. Examples are similar to interrupted attempts, except that the individual stops him/herself, instead of being stopped by something else. Has there been a time when you started to do something to make yourself not alive anymore (end your life or kill yourself) but you changed your mind (stopped yourself) before you actually did anything? What did you do? If yes, describe:		Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of aborted or self-interrupted _____	Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of aborted or self-interrupted _____
Preparatory Acts or Behavior: Acts or preparation towards imminently making a suicide attempt. This can include anything beyond a verbalization or thought, such as assembling a specific method (e.g., buying pills, purchasing a gun) or preparing for one's death by suicide (e.g., giving things away, writing a suicide note). Have you done anything to get ready to make yourself not alive anymore (to end your life or kill yourself)- like giving things away, writing a goodbye note, getting things you need to kill yourself? If yes, describe:		Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of preparatory acts _____	Yes No <input type="checkbox"/> <input type="checkbox"/> Total # of preparatory acts _____
		Most Recent Attempt Date:	Most Lethal Attempt Date:
		Initial/First Attempt Date:	
Actual Lethality/Medical Damage: 0. No physical damage or very minor physical damage (e.g., surface scratches). 1. Minor physical damage (e.g., lethargic speech; first-degree burns; mild bleeding; sprains). 2. Moderate physical damage; medical attention needed (e.g., conscious but sleepy, somewhat responsive; second-degree burns; bleeding of major vessel). 3. Moderately severe physical damage; <i>medical</i> hospitalization and likely intensive care required (e.g., comatose with reflexes intact; third-degree burns less than 20% of body; extensive blood loss but can recover; major fractures). 4. Severe physical damage; <i>medical</i> hospitalization with intensive care required (e.g., comatose without reflexes; third-degree burns over 20% of body; extensive blood loss with unstable vital signs; major damage to a vital area). 5. Death		Enter Code _____	Enter Code _____
Potential Lethality: Only Answer if Actual Lethality=0 Likely lethality of actual attempt if no medical damage (the following examples, while having no actual medical damage, had potential for very serious lethality: put gun in mouth and pulled the trigger but gun fails to fire so no medical damage; laying on train tracks with oncoming train but pulled away before run over). 0 = Behavior not likely to result in injury 1 = Behavior likely to result in injury but not likely to cause death 2 = Behavior likely to result in death despite available medical care		Enter Code _____	Enter Code _____